



African Journal of Biological Sciences



Qualitative Analysis of Phytochemicals of *Cassia tora* Linn. Seeds: A Review

Deepak Durgapal¹, Nitin Kumar Agrawal^{2*}, Animesh Agarwal², Manish Saxena², Pradeep Durgapal³ and Rashmi Singh¹

¹Department of Chemistry, Sardar Bhagat Singh Government P. G. College, Rudrapur, Uttarakhand, India

²Department of Applied Sciences and Humanities, Moradabad Institute of Technology, Moradabad, Uttar Pradesh, India

³Department of Chemistry, Government P. G. College, Bazpur, Uttarakhand, India

*Corresponding Author E-mail: nitinkumarag@yahoo.co.in

Abstract

Cassia tora Linn., an annual plant, is widely grown throughout Southeast Asia, including in India, Northern Australia, and the Americas. The enormous therapeutic powers of *Cassia* species are well known in the Indian medical system. The plant's leaves, seeds, and roots are among the parts that are said to have medicinal properties. It is widely used in traditional medicine as a laxative and is effective in treating leprosy, ringworm infection, ophthalmology, skin and liver issues, and diabetes mellitus. Several phytochemicals such as saponin, fixed oil/fats, protein, carbohydrates, tannins, flavonoids, alkaloids, gum, phenolic compounds, terpenoids, steroids, phenols, anthraquinones, glycosides, naphtho- α -pyrone toralactone, chrysophanol, physcion, emodin, rubrofusarin, and chrysophonic acid-9-anthrone have been reported. The presence of some phytochemicals such as tannin, saponin, and steroids, was demonstrated in the medicinal properties of the plant in its therapeutic applications by various researchers. This review article includes a literature review on the phytochemicals found in *Cassia tora* Linn. seeds.

Keywords: *Cassia tora*, phytochemical studies, solvent extract, seeds

Article History

Volume 6, Issue 5, 2024

Received: 01 May 2024

Accepted: 09 May 2024

doi: [10.33472/AFJBS.6.5.2024.2527-2533](https://doi.org/10.33472/AFJBS.6.5.2024.2527-2533)

Introduction

Plants are a major source of medications and are essential to maintaining global health (Constabel, 1990). In almost every civilization, both ancient and modern, plants have been employed as medicine (Deshpande and Bhalsing, 2013). Over 80% of the world's population, according to the WHO, relies on traditional medical practices, many of which are plant-based, to address their basic medical needs (Mazid et al., 2012). For all illnesses, the traditional medical system in our nation is crucial to the health of rural residents (Taid et al., 2014). Since the Rigveda and Atharvaveda, the curative potential of conventional herbal remedies has been recognised and documented (Bhattacharjya and Borah, 2008). Plants and their extracts have been utilised therapeutically ever since, and even today, medicines made from plants remain crucial to the global health care system (Yadav et al., 2006). About 45,000 plant species are found in India, and more than 35,000 of these are thought to have medicinal characteristics and are employed in various human societies around the globe (Lewington, 1993). Plants are a great source of a variety of bioactive substances that are used either directly or indirectly to treat a variety of human diseases (Dogra et al., 2015). Many cultures have employed plants from the genus *Cassia* in traditional medicine. 580 species of trees, shrubs, and plants belong to the genus *Cassia* (Malik et al., 2020).

The family Caesalpinaceae, which includes *Cassia tora* Linn, is primarily found in the tropics, West China, India, and Sri Lanka. It grows on dry soil throughout tropical regions, high hills with elevations up to 1800 m, as well as the plains in India, where it is a wasteland rainy season weed (Jain and Patil, 2010). The plant is 30 to 90 cm tall, with green leaves, yellow flowers, and subtetragonous obliquely septate pods that are 15 to 23 cm long and contain 23 to 30 seeds per pod [Fig. 1(a), Fig. 1(b) and Fig. 1(c)] (Kamble and Shubhangi, 2019). According to the conventional system of Indian medicine, this plant has a number of therapeutic characteristics (Pawar and D'mello, 2011). According to Ayurveda, leaves and seeds are astringent, laxative, antiperiodic, anthelmintic, ophthalmic, liver tonic, cardiotoxic, expectorant, etc., and are beneficial for treating leprosy, ringworm, colic, dyspepsia, constipation, cough, bronchitis, and cardiac problems (Bhot and Barua, 2015).



Fig. 1(a): Plant of *Cassia tora* Linn.



Fig. 1(b): Flowers of *Cassia tora* Linn.



Fig. 1(c): Seeds of *Cassia tora* Linn.

The seeds of *C. tora* have yielded a number of anthraquinones (Raghunathan et al., 1974). In Chinese medicine, the seeds of *C. tora* have been used as aperients, antiasthma treatments, diuretics, and to enhance visual function (Asolkar et al., 1992). The plant significantly contributes to a variety of biological functions such as antioxidant, hepatoprotective, hypolipidemic, antibacterial, antifertility, antimutagenic, antitumor, anti-inflammatory, antifungal, and antihelminthic activities due to various phytochemicals such as anthraquinone glycosides, naphthopyrone glycosides, phenolic compounds, flavonoids, sennosides, rubrofusarin triglucoside (Das et al., 2011). Therefore, the focus of this study work is on the qualitative analysis of phytochemicals found in *Cassia tora* Linn. seeds.

Phytochemical Analysis

The ethnopharmacological uses of medicinal plants in treatment, including the prevention of many diseases and disorders, have some biochemical support because of the phytochemical analyses of plants (Okigbo et al., 2009). After reviewing the literature, there are several phytochemicals, reported by various researchers, found in *Cassia tora* seeds: Saponin, fixed oil/fats, protein, carbohydrates, tannins, flavonoids, alkaloids, gum, and phenolic chemicals have been investigated (Supare and Patil, 2015). Alkaloids, flavonoids, and terpenoids were present, but saponins and tannins were not, (Sahadeo et al., 2014). The presence of saponins, tannins, flavonoids, and steroids was investigated (Khatak et al., 2014). In contrast to the absence of carbohydrates and gum, researchers observed the presence of alkaloids, flavonoids, terpenoids, saponins, tannins, phenols, amino acids/proteins, anthraquinones, glycosides, and steroids (Suradkar et al., 2017). The presence of tannin, saponin, protein, steroids, terpenoids, carbohydrates, alkaloids, flavonoids, and glycosides was studied (Shaikh and Syed, 2016). Alkaloids, flavonoids, terpenoids, carbohydrates, anthraquinones, glycosides, and the absence of saponins and tannins were all reported to be present (Patil and Shah, 2019). Steroids, carbohydrates, and proteins were present, but alkaloids and tannins were absent in the seeds of *C. tora*. (Pandya et al., 2017). In the investigation of seed mucilage, it was found that there were carbohydrates, proteins, and lipids present but no flavonoids, tannins, saponins, sterols, alkaloids, terpenoids, or starch (Singh et al., 2010). *C. tora* seeds contain quercetin and its

analogue, as well as emodin, chrysophanic acid, 1,8-dihydroxy anthroquinone, β -sitosterol, rhein like aglycones, cassiaside, rubrofusarin, and torosachryson (Yen et al., 1998).

The novel compounds torachryson tetraglucoside, nor-rubrofusarin gentiobioside, demethylflavasperone gentiobioside, torachryson gentiobioside, rubrofusarin triglucoside, and torachryson apioglucoside were isolated from the seeds of *Cassia tora* (Hatano et al., 1999). It was reported that two new naphtho-pyrone glycosides, 9-[(β -D-glucopyranosyl-(1 \rightarrow 6)-O- β -D-glucopyranosyl)oxy]-10-hydroxy-7-methoxy-3-methyl-1H-naphtho[2,3-c]pyran-1-one and 6-[(α -apiofuranosyl-(1 \rightarrow 6)-O- β -D-glucopyranosyl)oxy]-rubrofusarin, together with cassiaside and rubrofusarin-6- β -gentiobioside were also discovered from the seeds of *Cassia tora* Linn. (Wong et al., 1989).

The seeds of *C. tora* contain sitosterol in petroleum ether extract. Chrysophanol, physcion, emodin, and rubrofusarin were present in CHCl_3 extract, and ethanolic extract also shows glycosides II and I, in which Nitrofurarin-6- β -gentiobioside was recognised as glycoside II (Jain and Patil, 2010). The hydroalcoholic seed extracts of *Cassia tora* showed high anti-diabetic effects (Adamu and Oladosu, 2023).

Naphtho- α -pyrone toralactone, chrysophanol, physcion, emodin, rubrofusarin, and chrysophonic acid-9-anthrone were all present in the seeds of *Cassia tora*. From the butanol soluble extract of the seeds, three naphthopyrone glucosides were isolated: cassiaside, rubrofusarin-6-O- β -D-gentiobioside, and toralactone-9-O- β -D-gentiobioside (Soumyanath, 2005; Mukherjee, 2002).

Conclusion

Cassia tora is one of the most significant sources of medicinally useful phytochemicals. It is widely used in both Chinese and Ayurvedic medicine. The contents of the phytochemicals in *Cassia tora* seeds were covered entirely in the information that was compiled from contemporary literature sources. Among the medicinally useful phytochemicals, alkaloids, carbohydrates, saponins, glycosides, proteins, tannins, phenols, anthraquinones, terpenoids, and other bioactive compounds are most significant, but the chemical components vary qualitatively and quantitatively not only from plant to plant but also in different samples of the same species depending on various atmospheric factors and storage conditions. As a result, it can be said that *Cassia tora* Linn. contains a wide range of phytochemicals that are useful in different ways to treat a wide range of disorders. As a result, *Cassia tora* has a lot of potential for research and may be further utilised as a source of beneficial phytochemical compounds for the pharmaceutical industry.

References

Adamu, M. W. and Oladosu, J. I. (2023). Phytochemical screening and anti-diabetic effects of the hydroalcoholic seed and leaf extracts of Pot Cassia (*Cassia tora*). *Journal of Clinical and Metabolism Studies*, 2(2), 31-40.

Deepak Durgapal/ Afr.J.Bio.Sc. 6(5) (2024) 2527-2533

Asolkar, L.V., Kakkar, K. K. and Chakre, O. J. (1992). Second supplement to glossary of Indian medicinal Plants, PID, CSIR, New Delhi, 180-181.

Bhattacharjya, D. K. and Borah, P. C. (2008). Medicinal weeds of crop fields and role of women in rural health and hygiene in Nalbari district, Assam. *Indian J. Tradit. Knowl.* 7(3), 501-504.

Bhot, M. and Barua, M. (2015). Pharmacognostical, Physico-chemical and Phytochemical Evaluation of leaves of *Cassia tora* and *Cassia fistula*. *Int. J. of Life Sciences*, 4, 1-11.

Constabel, F. (1990). Medicinal plant biotechnology. *Planta Med.*, 56(05), 421-425.

Das, C., Dash S., Sahoo D. C., Mohanty A. and Rout D. (2011). *Cassia tora*: A Phyto-Pharmacological Overview. *Int. J. Res. Ayurv. Pharm.*, 2(4): 1162 - 1174.

Deshpande, H. A. and Bhalsing, S. R. (2013). Recent advances in the phytochemistry of some medicinally important *Cassia* species: A review. *Int. J. Pharm. Biol. Sci.*, 2(3), 60-78.

Dogra, K. S., Chauhan, S. and Jalal, J. S. (2015). Assessment of Indian medicinal plants for the treatment of asthma. *J. Med. Plant Res.*, 9(32), 851-862.

Hatano, T., Uebayashi, H., Ito, H., Shiota, S., Tsuchiya, T. and Yoshida, T. (1999). Phenolic constituents of *Cassia* seeds and antibacterial effect of some naphthalenes and anthraquinones on methicillin-resistant *Staphylococcus aureus*. *Chemical and Pharmaceutical Bulletin*, 47(8), 1121-1127.

Jain, S. and Patil, U. K. (2010). Phytochemical and pharmacological profile of *Cassia tora* Linn.-An Overview. *Indian Journal of Natural Products and Resources*, 1(4), 430-437.

Kamble, K. J. and Shubhangi, D. (2019). Study of physico-chemical properties of *Cassia tora* Linn. leaves powder. *International Journal of Agricultural Engineering*, 12(2), 235-242.

Khatak, S., Sharma, P., Laller, S. and Malik, D. K. (2014). Antimicrobial, antioxidant and phytochemical property of *Cassia tora* against pathogenic microorganisms. *Journal of Pharmacy Research*, 8(9), 1279-1284.

Lewington, A. (1993). Medicinal plants and plant extracts: a review of their importation into Europe. *Traffic International, Cambridge, UK*.

Malik, J. K., Yadav, B., Yadav, A. P. and Soni, H. (2020). Phytography & Phytopharmacology of Genus *cassia*. *Journal of Advances in Bio-pharmaceutics and Pharmacovigilance*, 2(2), 1-9.

Mazid, M., Khan, T. A. and Mohammad, F. (2012). Medicinal plants of rural India: a review of use by Indian folks. *Indo-Glob. Res. J. Pharm. Sci.*, 2(3), 286-304.

Deepak Durgapal/ Afr.J.Bio.Sc. 6(5) (2024) 2527-2533

Mukherjee, P. K. (2002). Quality Control of Herbal Drug (1st ed., pp. 531, 554–557, 714–716). Business Horizons, New Delhi.

Okigbo, R. N., Anuagasi, C. L. and Amadi, J. E. (2009). Advances in selected medicinal and aromatic plants indigenous to Africa. *Journal of Medicinal Plants Research*, 3(2), 86-95.

Pandya, M. P., Sameja, K. D., Patel, D. N. and Bhatt, K. D. (2017). Antimicrobial activity and phytochemical analysis of medicinal plant *Cassia tora*. *International Journal of Pharmacy and Chemistry*, 3(4), 56-61.

Patil, S. and Shah, R. (2019). Standardization, formulation & antimicrobial activity of *Cassia tora*. *Int. J. Pharm. Sci. and Res.*, 10(4), 1870-1879.

Pawar, H. A. and D'mello, P. M. (2011). *Cassia tora* Linn.: an overview. *International Journal of Pharmaceutical Sciences and Research*, 2(9), 2286-2291.

Raghunathan K., Hariharan V. and Rangaswami S. (1974). Chrysophanol-1- β -gentiobioside, a new anthraquinone glycoside from *Cassia tora* Linn. *Indian J. Chem.* 12, 1251-1253.

Sahadeo, P., Wankhade, S., Somkuwar, D. and Kamble, V. (2014). Phytochemical and chromatographic fingerprint studies on chloroformic extracts of *Cassia tora* L. *Int. J. Drug Dev. and Res*, 6(3), 0975-9344.

Shaikh, R. and Syed, I. Z. (2016). Phytochemical, proximate and nutrient analysis of *Cassia Tora* seeds. *International Journal of Pharmaceutical Science Invention*, 5(6), 04-06.

Singh, S., Bothara, D. S. B., Singh, S., Patel, R. D. and Mahobia, N. K. (2010). Pharmaceutical characterization of *Cassia tora* of seed mucilage in tablet formulations. *Scholars Research Library*, 2(5), 54-61.

Soumyanath, A. (2005). Traditional Medicines for Modern Times (1st ed., p. 11). Taylor and Francis, USA.

Supare, S. and Patil, M. (2015). Estimation of phytochemical components from *Cassia tora* and to study its larvicidal activity. *International Journal of Pharmaceutical Science Invention*, 4(6), 11-16.

Suradkar, V. B., Wankhade, B. B. and Dabbe, P. G. (2017). Phytochemical analysis of some contents of *Cassia tora* and *Xanthium strumarium* plant seeds. *International Journal of Advanced Research in Science, Engineering and Technology*, 4(4), 3727-3731.

Taid, T. C., Rajkhowa, R. C. and Kalita, J. C. (2014). A study on the medicinal plants used by the local traditional healers of Dhemaji district, Assam, India for curing reproductive health related disorders. *Adv. Appl. Sci. Res.*, 5(1), 296-301.

Deepak Durgapal/ Afr.J.Bio.Sc. 6(5) (2024) 2527-2533

Wong, S. M., Wong, M. M., Seligmann, O. and Wagner, H. (1989). New antihepatotoxic naphtho-pyrone glycosides from the seeds of *Cassia tora*. *Planta Medica*, 55(3), 276-280.

Yadav, J. P., Kumar, S. and Siwach, P. (2006). Folk medicine used in gynecological and other related problems by rural population of Haryana. *Indian J. Tradit. Knowl.* 5(3), 323-326.

Yen, G. C., Chen, H. W. and Duh, P. D. (1998). Extraction and identification of an antioxidative component from Jue Ming Zi (*Cassia tora* L.). *Journal of Agricultural and Food Chemistry*, 46(3), 820-824.